

# Global Surface Temperatures of the Moon

Jean-Pierre Williams (UCLA)

David A. Paige (UCLA)

NASA Exploration Science Forum 2014

UCLA JPL




STONY  
BROOK




# Overview



- What Diviner does:
  - Measure reflected solar and emitted infrared radiation.
  - Nine spectral channels with wavelengths ranging from 0.3 to 400 microns.
- Five years of data → global diurnal temperatures can be adequately resolved to create global maps of surface temperatures.
- Bolometric brightness temperatures from radiance from the 7 infrared spectral channels
  - measure of the spectrally integrated flux of infrared radiation emerging from the surface (*Paige et al., 2010*).



- Sub-spacecraft longitude cycles between  $-180^\circ$  and  $180^\circ$  every  $\sim 27$  days
- Sub-spacecraft local time cycles between 0 and 24 hours once every Earth year
- Daytime and nighttime observations on alternate hemispheres in a single orbit
  - two different local times separated by  $180^\circ$  longitude.




Dawn







←  
time

First light




# Diviner Global Diurnal Bolometric Temperature Coverage




Rocks, Regolith Thermophysical Properties and Vertical Structure

# Diviner Global Coverage at 64ppd: 5:45 AM – 6:15 AM


ESMD Mission: 16%




SMD Mission: 32%




ESM1 Mission: 27%




Combined: 55%





July 2009 - March 2012






July 2009 – June 2014












Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature






Bolometric Temperature




Bolometric Temperature







Bolometric Temperature






Bolometric Temperature













Bolometric Temperature




Bolometric Temperature




Bolometric Temperature




Bolometric Temperature







# Maximum Global Temperatures





# Minimum Global Temperatures




# Minimum Global Temperatures









Local time: 20 hr




Local time: 21 hr




Local time: 22 hr




Local time: 23 hr




Local time: 24 hr




Local time: 1 hr



Local time: 2 hr




Local time: 3 hr




Local time: 4 hr

## Nighttime $T_{bol}$ – Nighttime Averaged $T_{bol}$



# Summary

- Systematic observations of the global thermal state of the Moon
- Mapping global diurnal variability
- Characterize the surface energy balance
  - How does the lunar regolith stores and exchanges heat.
- Extremely complex thermal environment
  - Highly insulating surface
  - Lack of an appreciable atmosphere to buffer surface temperatures
  - Slow rotation
- Terminators and at high latitudes topography dominates the surface temperatures.
- Hottest temperatures, which are in near-equilibrium with the solar flux, are influenced by radiative properties of the surface.
- Nighttime temperatures determined by the radiation of sensible heat stored in the subsurface during the day and are sensitive to the thermophysical properties of the regolith.




[jpierre@mars.ucla.edu](mailto:jpierre@mars.ucla.edu)



# Diviner Spectral Channels

Diviner Spectral Channels' Passbands



Channel Number	Channel Type	Channel Name	Passband $\mu\text{m}$	Measurement Function	Minimum Detectable Temperature (K)
1	Solar	High Sensitivity Solar	0.35-2.8	Reflected solar radiation, high sensitivity	-
2	Solar	Reduced Sensitivity Solar	0.35-2.8	Reflected solar radiation, reduced sensitivity	-
3	8 $\mu\text{m}$	7.8 $\mu\text{m}$	7.55-8.05	Christiansen feature	150
4	8 $\mu\text{m}$	8.25 $\mu\text{m}$	8.10-8.40	Christiansen feature	150
5	8 $\mu\text{m}$	8.55 $\mu\text{m}$	8.38-8.68	Christiansen feature	150
6	Thermal	13-23 $\mu\text{m}$	13-23	Surface Temperature (most sensitive channel for >178 K)	856
7	Thermal	25-41 $\mu\text{m}$	25-41	Surface Temperature (most sensitive channel for 69-178K)	55
8	Thermal	50-100 $\mu\text{m}$	50-100	Surface Temperature (most sensitive channel for 43-69 K)	40
9	Thermal	100-400 $\mu\text{m}$	100-400	Surface Temperature (most sensitive channel for <43 K)	30

# Bolometric Brightness Temperature (TBOL)

- Measure of spectrally integrated flux of infrared radiation from the surface
- TBOL more directly related to heat balance of the surface
- Individual Diviner channels can exhibit anisothermality

In radiative equilibrium, the surface heat balance can be expressed as:

$$\sigma T_{BOL}^4 = \pi \int_0^{\infty} \varepsilon(\lambda) B(\lambda, T_s) = S_0(1 - A)$$
$$\sigma T_{BOL}^4 = \sum_{i=3}^9 \sigma T_i^4 f(T_i, \lambda_1, \lambda_2) \quad f(T_i, \lambda_1, \lambda_2) = \frac{\int_{\lambda_1}^{\lambda_2} B(\lambda, T) d\lambda}{\int_0^{\infty} B(\lambda, T) d\lambda}$$

For detailed description see:

Paige et al., *Diviner Lunar Radiometer Observations of Cold Traps in the Moon's South Polar Region, Science, Vol 330, p479-482, 2010*